

Index

References to tables and figures are given as, for example, 98*t*, 306*f*.

- abduction splints, 224, 245, 263, 350
- ABILHAND-Kids, 140
- accessibility, 391–92
- active joint range (range of movement), 131–33
 - see also* joints
- activities of daily living (ADL) *see* daily life
- activity, definition, 4, 5
- activity limitations, definition, 5
- acupuncture, 45
- adduction, 297–98
- adolescents, 2, 9–10, 11, 61–66, 67–70, 235, 236, 249, 344, 353
 - relationship with therapist, 65–66
 - surgery, 366
- adults, 2, 10, 11, 249, 342, 353
- aetiology, 7
- afferent stimuli, 74, 77–78, 118–19
- Alberta Infant Motor Scale (AIMS), 145
- alternative treatments, 44–46
- American Academy for Cerebral Palsy and Developmental Medicine (AACPD), 76, 81, 97, 98*t*, 359
- anarthria, 311
- ankle-foot orthoses, 68*f*, 250, 251, 259, 264–66, 345, 347, 363
- anteroposterior weight shift (rocking), 169, 232, 247*f*, 249, 252*f*
- anticipatory grasp, 298, 302
- antigravity mechanism, 21, 106, 128, 230, 233*f*, 237, 245, 341, 346
- arms *see also* shoulder
 - assessment of, 123–24, 128, 137–38
 - atypical posture, 242, 346
 - deformities, 344–45, 346, 353, 358–59, 365
 - developmental levels, 115, 169–170, 221–22
 - developmental training, 176–77, 181–82, 184, 187–89, 190–91, 205, 216*f*, 252–53
 - orthoses (splints, gaiters), 305–8
 - postural control, 109–10, 216*f*, 278–282
 - propping, 184
- aromatherapy, 45
- arthritis, 64, 66
- Ashworth Scale, 129
- Asian families, 30
- assessment, 2, 110, 121–156, 161
 - attention span, 123
 - atypical posture, 19, 159, 220–23, 225, 227, 238–242
 - changes in function, 9–10, 62–63, 66–67
 - combined assessments, 43, 324
 - equipment, 154–55, 371–386
 - evaluation of research studies, 94–95, 96–102
 - functional measures, 324–25
 - home and school visits, 30, 121–22, 160, 324
 - measurement tools, 125–145, 233–36
 - endurance, 127, 128–29, 136
 - functional measures, 133–141
 - gait analysis, 145–153*t*, 263, 344, 345–48
 - neonatal assessment, 144–45
 - norm-referenced measures, 143–44
 - quality of life (QOL), 142–43
 - ranges of motion, 131–33
 - selective motor control, 126–27
 - strength, 127–29
 - timed tests, 128–29, 136
 - wheelchair skills, 136, 148, 386–88, 397
 - motor delay, 22–23
 - neuroimaging, 7–8
 - observations, 123–24
 - of outcomes, 122–23
 - pain, 132
 - parent-child interaction, 40–41
 - preterm infants, 10
 - prognosis for walking, 62–63, 235–36
 - purpose, 122–23
 - record-keeping, 155–56
 - reflexes, 21, 74, 76, 78, 149*t*–153*t*, 235, 270, 315
 - task analysis, 32–34, 55–56, 80, 88–89, 121–22, 392
 - and therapy planning, 159–160
 - 'traffic light' evaluation, 77
- Assessment of Motor and Process Skills (AMPS), 139–140

Treatment of Cerebral Palsy and Motor Delay, Sixth Edition. Sophie Levitt and Anne Addison.

© 2019 Sophie Levitt and Anne Addison. Published 2019 by John Wiley & Sons Ltd.

- Assisting Hand Assessment (AHA), 137
 assistive technology *see* equipment
 associated impairments and disabilities, 3–4, 18, 20, 21, 22–23, 161–68, 299–300, 311, 333, 386–87, 390–91
 Association of Paediatric Chartered Physiotherapists, 309, 357, 359, 360
 asymmetrical tonic neck reflex (ATNR), 150*t*, 202–4, 341
 asymmetry, 166–67, 229, 240, 244, 248, 255, 340–42, 352
 ataxia, 248
 ataxic cerebral palsy, 9, 20–21, 237
 athetosis, 18–19, 73, 205, 248, 298, 342, 353 *see also* dyskinesia
 'athetoid dance', 20, 64, 237
 fluctuating tone, 117, 338
 attention, development of, 52–54, 123, 330
 atypical (stereotyped) synergies, 340
 atypical development, 1–2, 3, 8–11, 15–17, 22–23, 158
 gait, 19, 20
 reflexes, 21–22
 synergies, 72, 119–120, 127, 341
 atypical postures, 238–242, 244–45, 263–68
 asymmetrical postural stabilisation, 240
 correction of, 253
 unsupported standing, 241
 well-supported standing, 242
 auditory development, 164, 286–87
 augmentative/alternative communication (AAC), 390–91
 autism, 3
 avoiding reactions, 302
 Ayres, A.J., 81
- babies (infants, neonates), 10–11, 106
 hypotonicity, 117, 200
 neonatal assessment, 144–45
 preterm (premature) babies, 10–11
 reflexes, 21–22, 197–98, 201–2, 204, 227, 235, 237, 287, 315, 344
 back, 128
 baclofen, 17, 87, 116, 266
 balance *see* postural control
 bathing, 30, 368–69, 378
 Bayley Scales of Infant Development, 143
 bear-walk, 170, 172*f*, 189, 190–91
 behaviour, 38, 319–320, 321–22
 behavioural impairments, 3, 52, 53, 123, 160, 332
 definition, 51–52
 in feeding, 321–22
 in group work, 331–32
 and personality, 5–6, 20, 333
- Berg Balance Scale, 136
 bilateral cerebral palsy, 14–15, 16*f*, 18, 54*f*, 62–63, 124, 222, 267, 316–17, 339–340, 346–48, 357–58
 Bimanual Fine Motor Function (BFMF)
 classification, 13
 bimanual therapy, 285–86
 biofeedback, 58, 242
 biomechanics, 64, 83, 112–13, 239, 241, 248, 255, 263, 265, 342–43, 345
 'bird-wing' position, 281*f*
 blindness, 161
 Bobath, B., 75–77, 111
 Bobath, K., 75–77, 111
 Bobath approach (NDT), 75–77
 body functions and structures,
 definition, 4, 5
 body harnesses, 84–85, 378
 body image development, 165, 352
 bone density, 236, 379–380
 boots (shoes, trainers), 265–66, 363
 bottom shuffling, 124, 167, 174, 192
 botulinum toxin A, 116, 250, 295, 307, 340, 345, 347, 353, 354, 357–361
 Bower, E., 95
 bow legs (varus knees), 339
 braces, 72, 363 *see also* orthoses
 brain, 7–8, 76
 neuroplasticity, 89–90
 traumatic brain injury, 7, 160
 Brazelton, T., 47–48
 breastfeeding, 149*t*, 278
 breathing problems, 18, 65, 68, 114, 173, 366
 bridging (hips), 118, 205
 British Assistive Technology Association, 391
 British Institute for Brain Injured Children (bibic), 74
 built environment, accessibility, 391–92
 'bunny hopping', 55, 124, 192
- calipers, 72, 256*f*
 Canadian Occupational Performance Measure (COPM), 140–41
 cardinal points, 149*t*
 carers, 28, 63, 389–390
 carrying the child, techniques, 204, 224, 225, 307*f*, 308*f*
 case-control studies, 97
 case studies, 95
 casts (plaster casts), 99, 245, 264–65, 351, 352–57, 359, 362–63
 cerebral palsy, 1–24 *see also* specific types, e.g. ataxic cerebral palsy, choreoathetotic cerebral palsy, dyskinesia, dystonia, spasticity (spastic hypertonus)

- adolescents with cerebral palsy, 2, 9–10, 11, 61–66, 67–70, 236, 344, 353, 366
 adults with cerebral palsy, 2, 10, 11, 236, 342, 353
 associated impairments and disabilities, 3–4, 18, 20, 21, 22–23, 161–68, 299–300, 311, 333, 386–87, 390–91
 causes, 7, 160
 classification systems *see* classification systems
 definition, 4–6, 7
 incidence, 1, 7
 onset, 160
 Cerebral Palsy Integrated Pathway (CPIP-UK), 131–32
 cerebral palsy therapists, 75
 Chailey Levels of Ability, 136
 chairs/seating systems, 205, 215, 217–19, 221, 223–25, 315, 369, 370, 371–79
 adjustments to, 374
 developmental stages, 217–218
 evaluation and research, 372–373
 Matrix seating, 378, 379*f*
 measurement, 371–372*f*
 powered mobility aids, 383–84, 386–88
 tilt-in-space seating, 375–76*f*, 387*f*, 388
 wheelchairs, 62, 66, 68–69, 369, 378, 383–84, 385
 wheelchair skills, 136, 148, 386–88, 397
 Cheyne Centre for Children with Cerebral Palsy, 330
 Children's Assessment of Participation and Enjoyment (CAPE), 140
 Children's Hand-use Experience Questionnaire (CHEQ), 140
 choking, 317, 319–320, 321
 choreoathetotic cerebral palsy, 19, 20
 classification systems, 11–21
 Bimanual Fine Motor Function (BFMF)
 classification, 13
 Communication Function Classification System (CFCS), 13–14
 Eating and Drinking Ability Classification System (EDACS), 14, 315
 Gross Motor Function Classification System (GMFCS), 11–12, 14, 62, 95, 124
 Manual Ability Classification System (MACS), 12–13, 14
 topographical classifications, 3, 14–15
 clenched toes, 241
 Cochrane reviews, 85, 97, 312, 381–82
 co-contraction, 18, 116, 339
 atypical, 128, 230, 338, 340–41, 346
 typical, 78, 106, 126
 cohort studies, 97
 collaborative learning approach, 30–31, 61–62, 315
 contracts, 26
 Collis, E., 75
 communication, 6, 158, 166, 310–313
 AAC devices, 390–91
 with adolescents and adults, 62, 64, 65–66, 69–70
 assessment of, 123
 clients with learning disabilities, 160
 clients with visual impairments, 161–62, 166–67
 cross-cultural communication, 29–30
 cues for learning, 56–58, 164–68
 developmental levels, 310–11
 and dyskinesia, 20
 face-to-face contact, 313
 facial expressions, 14, 26, 32, 310, 390
 feedback, 56–58, 75, 77, 88
 gesture, 14, 32, 123, 137, 278, 310
 non-verbal communication, 3, 32, 52, 124, 160, 311, 313
 parent-child interaction, 37, 38–41, 46–48, 163–64, 166–67, 312–13
 practical suggestions, 312–13
 sign language, 124, 311–12
 speech and language impairments, 311–13
 Communication Function Classification System (CFCS), 13–14
 community therapy, 324
 complementary treatments, 44–46
 conductive education, 77, 79–81, 330
 confidence interval, definition, 97
 confidentiality, 61, 68, 102
 constraint-induced movement therapy, 6, 282–85
 context therapy, 81
 contractures *see* deformities
 control group, definition, 98
 Cotton, E., 111, 119
 counselling, 27, 41, 61, 68, 69
 counterpoising (postural adjustment), 106, 137, 168*f*, 169–170, 187–88, 190, 193–94, 208, 209, 227*f*, 231–32, 241, 252–55, 276–77, 279*f*
 counter rotations, 198
 CPIP-UK (Cerebral Palsy Integrated Pathway), 131–32
 Craighalbert Centre, 81
 cranial osteopathy, 45
 crawling, 55, 111, 114–15, 124, 167, 170, 180, 185, 192, 341
 creeping, 17–18, 78–79, 119, 124, 178–180, 183–85, 187–89, 204
 Crothers, B., 94
 crouch gait, 255, 340
 cruising, 170, 232, 234*f*, 246, 250, 251, 255–56, 327
 cryotherapy, 74, 351
 culturally appropriate therapy, 7, 9, 29–30, 40, 46, 155, 174

- daily life, 2, 5–7, 62, 63, 309–28
 assessment of participation, 138–141
 carrying the child, 224, 225, 307f, 308f
 child's own strategies, 50, 54–55, 206, 207f
 communication *see* communication
 conductive education, 79
 education, 318, 324–26, 383
 environmental factors, 5, 30, 32, 160, 318, 344, 383, 385–86, 388, 391–92
 feeding, 18, 22–23, 62, 163, 278, 289, 314–322, 369–370, 375
 motor learning (coaching), 50–54, 59–60, 158–59, 183–84, 300, 360
 movement in bed, 63–64, 115, 183–84, 267
 parent-child interaction, 37, 38–41, 46–48, 163–64, 166–67, 312–13
 play, 6, 36–37, 54, 163, 165, 166–67, 182, 230, 289, 326–28, 329–330, 334–35
 positioning, 159, 204, 238, 314, 317–18, 349–352, 361, 363, 370, 388–390
 quality of life (QOL), 62, 68–69, 96, 142–43, 326, 340
 self-care tasks, 62, 63, 69, 118, 160, 163, 164f, 313–324, 335, 368–370, 378
 sleep, 20, 22, 41, 265, 351–52, 366, 381–83
 sport and leisure activities, 67–69, 70, 85–86, 230, 277, 326–28, 360
 task analysis, 32–34, 55–56, 121–22, 392
 Day, S.M., 62
 Debusse, D., 136, 141
 deformities, 16, 17, 89, 113–14, 160, 337–366
 in adolescents and adults, 64, 342, 344, 353, 366
 assessment of, 124, 131–33, 154, 337–38
 causes, 338–343, 376–77
 dynamic deformities, 16, 54–55, 244–45, 337, 340, 352–57, 361–66
 'frog' position, 174, 182, 184, 203, 338–39
 kyphosis, 210f, 221, 222, 223–24, 281f, 314, 316, 347, 366, 372, 378
 lordosis, 177, 192, 205, 208, 246, 260f, 347, 354f, 355f, 361, 375
 prevention, 54–55, 80, 191, 192–93, 215, 222, 223–24, 245, 256f, 257f, 266, 307
 scoliosis, 65, 120, 194, 203f, 222, 223, 224, 226f, 240, 341, 342, 343, 347, 350, 355f, 365–66, 371, 378, 381–82
 torticollis, 204, 222, 341
 windswept posture, 201, 203f, 229–230, 341, 350–51, 365–66, 377
 Denver Developmental Screening Test (DDST), 144
 Department of Health (UK), 144
 developmental levels, 75–77, 89, 111–16, 157–168, 193–208, 220–233, 237–263, 393–97
 0–3 months, 171f, 173–180, 200–204, 220–21, 233f, 237, 286–87, 292, 310, 315, 393–396
 0–6 months, 195f, 196–98, 237–38, 327
 3–5 months, 287–89, 292
 3–6 months, 171f, 180–85, 221–25, 233f, 310, 315, 393–396
 4–6 months, 204–5
 5–7 months, 233f, 289–300
 6–9 months, 171f–72f, 181, 185, 195f, 206–8, 225–26, 310, 315–16, 393–396
 6–10 months, 198–200
 6–12 months, 322, 327
 7–9 months, 234f, 300
 9–12 months, 172f, 189, 195f, 226–230, 234f–35f, 246–48, 292, 300–301, 310, 316, 393–396
 12–18 months, 235f, 248–250, 292, 310, 316, 322, 325, 327, 393, 396
 18–24 months, 269–271, 292, 310, 316, 322, 325, 327, 396
 2–7 years, 272–77, 310, 316, 323, 325, 327–28, 396
 atypical development, 1–2, 3, 8–11, 15–17, 22–23, 157–58, 161, 341
 hand function, 280–82, 286, 287, 289–290, 295–305
 language acquisition, 311–13
 prone development, 173–74, 180–81, 189
 sitting, 220–26
 standing and walking, 237, 238–242, 246–47, 248–250, 271, 340, 349
 supine development, 200–204, 205, 206–8
 and biomechanics, 112–13, 238–39, 241, 248
 typical development, 9, 30, 53, 78, 83, 108, 114–16, 143–44
 cultural/individual variation, 7, 9, 29–30, 114
 hand function, 283t–85t, 286, 298, 322–23, 325
 language acquisition, 310–11
 prone development, 173, 174, 189, 393–94
 sitting, 220, 221, 225, 226–27, 395–96
 standing and walking, 237, 246, 248, 265–66, 269–271, 396
 supine development, 196–200, 204–5, 206, 394–95
 use of early levels by adolescents and adults, 63–64
 developmental training, 111–16
 hand function, 115, 166–67, 204, 214–15, 246, 252–53, 277–308, 324–26
 hand-eye coordination, 286–298, 300, 303, 304–5
 prone development, 168–193, 204
 sitting, 208–30
 standing and walking, 230–263, 267–277, 363, 364f
 supine development, 193–208
 diagnosis *see* assessment
Diagnostic and Statistical Manual of Mental Health Disorders (DSM-V), 22
 digital assistive technology, 390–91
 diplegia, 8, 15, 54f, 62–63, 124, 271, 316–17, 339–340, 346–48, 357–58 *see also* bilateral cerebral palsy

- disability, definition, 4
- DISABKIDS (DCGM-37), 142
- distractibility, 3, 53
- Doman-Delacato system, 73
- dorsal rhizotomy, 17, 87, 95–96, 116
- dorsiflexion, 241, 250, 255, 265, 340, 355–56, 363
- dressing, 62, 118, 164*f*, 335
- developmental levels, 322–23
- practical suggestions, 323–24
- drugs, 17, 86–87, 116 *see also* baclofen; botulinum toxin A
- dynamical systems theory, xv, 88, 89, 90, 92
- dynamic deformities, 16, 54–55, 244, 337, 340, 352–57, 361–66
- dysarthria, 311
- dyskinesia, 18–20, 124, 192–93, 205, 214–15, 222, 238–39, 248, 249, 281–82, 302, 341, 342, 353, 378
- fluctuating tone, 117, 338
- dysmetria, 20
- dysphagia, 67, 315, 317
- dyspraxia, 3
- dystonia, 17, 18–20, 192–93, 214–15, 298, 338, 339, 376
- Early Activity Scale for Endurance (EASE), 129
- Eating and Drinking Ability Classification System (EDACS), 14, 315
- echolalia, 310
- eclectic approach, xi, 2, 76–77, 103–20
- Edinburgh Visual Gait Analysis, 147
- education, 318, 324–26, 383
- elbow, 128, 133, 256*f*, 257*f*, 261
- electrical stimulation (FES/NMES), 83–84
- emotion
- and dyskinesia, 19–20, 278
- emotional development, 3–4, 22, 70, 163, 165, 318
- emotional support during therapy, 26–27, 31–32, 315, 318, 351
- and learning, 50–51, 52, 60, 79
- endurance, 127, 128–29, 136
- environmental factors, 5, 30, 32, 160, 318, 344, 383, 385–86, 388, 391–92
- epilepsy, 3, 18
- Equality Act (UK, 2010), 64
- equilibrium reactions, 107–8, 111
- equinus feet, 192, 223, 241, 244, 248, 337, 341, 342, 343, 345–47, 352, 357, 359, 362–63, 364
- equipment, 50, 56, 115, 118, 297, 367–392
- AAC devices, 390–91
- assessment of, 154–55, 371–386
- bathing and showering, 30, 368–69, 378
- body harnesses, 84–85, 378
- braces and calipers, 72
- and the built environment, 391–92
- and cultural context, 30
- digital assistive technology, 390–91
- lying, 202–4, 360
- for moving and handling, 63, 86, 267, 269, 369, 384–85, 388–390
- orthoses *see* orthoses
- play, 326, 328
- powered mobility aids, 383–84, 386–88
- prone standers, 238, 245
- regular review of, 368, 388
- seating systems, 205, 215, 217–19, 221, 223–25, 315, 369, 370, 371–79
- sleep systems, 351–52, 360, 366, 381–83
- standing frames, 50*f*, 73, 114, 237, 245–46, 379–381
- strength training, 82–83
- tables, 321, 371
- toilet seats, 369
- treadmills, 84–85
- tricycles, 192, 205, 228, 263, 277
- walking aids, 115, 168, 236–37, 256*f*, 257*f*, 258–263, 383, 384–85
- wedges, 176–78, 205, 224, 225, 350
- wheelchairs, 62, 66, 68–69, 369, 378, 383–84, 385
- wheelchair skills, 136, 148, 386–88, 397
- evaluation of treatment, 6–7, 28–29, 37–38, 71–89, 93–102, 309, 360–61
- everted toes, 364
- evidence-based practice, xiv, 10, 29, 93–102, 104–5, 309
- levels of evidence, 97–98
- exercise, importance of, 67–68
- extension, 200–201, 202, 203*f*, 213, 222
- extensor thrust, 21, 150*t*, 166, 202, 221, 268, 321, 341, 373, 375
- eyes
- dyskinesia, 20
- hand-eye coordination, 283*t*–85*t*, 286–298, 300, 303, 304–5
- face-to-face contact, 313
- facial expressions, 14, 26, 32, 310, 390
- failure, attitudes to, 50–51, 52, 60, 75
- falling (saving reactions), 107, 128, 153*t*, 170–71, 184, 191, 194, 210, 211*f*, 233, 240–41, 249–250, 270, 280
- families *see also* parents
- assessment of needs, 155
- cultural context, 7, 9, 29–30, 40, 46, 155, 174
- family-centred services, xii, 25–30, 46–48
- relationships with adolescents, 65
- sibling relationships, 43, 65
- and social/emotional development, 3–4, 22, 60, 163, 165, 318–320, 368–69
- support for parents, 26–28, 41–42, 315, 318, 348, 351

- fathers, 16*f*, 28, 35*f*, 36*f* *see also* parents
- fatigue, 19, 22, 52, 63, 66–67, 116
- Fay, T., 73
- feedback, 56–58, 75, 77, 88, 242
- feeding, 18, 22–23, 163, 278, 289, 314–322, 369–370, 375
- arm function, 314, 318–19, 321
 - developmental stages, 315–16
 - equipment, 320, 369–370
 - gastrostomy, 173, 315, 320
 - parent-child relationships, 318–320
 - practical suggestions, 316–18
 - specific difficulties, 320–22
- feet
- assessment of function, 133
 - deformities, 223, 250, 265–66, 345–46, 351, 362–64
 - developmental training, 191, 215, 222, 251–52
 - footwear, 265–66, 363
 - orthoses, 68*f*, 250, 251, 259, 264–66, 345, 347, 351, 363
 - reflexes, 149*t*, 150*t*
- fingers, 296–98, 302–3, 320
- Finnie, N., 77
- Five Times Sit to Stand (FTSS), 128–29, 136
- fixed deformities *see* deformities
- flexion, 119, 340
- dorsiflexion, 241, 250, 255, 265, 340, 355–56, 363
 - in grasping, 296–97
 - in prone development, 173–74, 186, 192
 - in sitting, 210*f*, 213, 215, 222, 224
 - in standing and walking, 239–240, 241, 250
 - in supine development, 200–201, 202, 206
- floor seats, 155, 224, 225, 361, 374, 378–79
- focus groups, definition, 102
- Foley, J., 249
- footwear, 265–66, 363
- four-point kneeling, 153*t*, 361
- 'frog' position, 174, 182, 184, 200, 201*f*, 203, 338–39
- functional electrical stimulation (FES), 83
- Functional Independence Measure for Children (WeeFIM), 139
- Functional Mobility Scale (FMS), 139
- functioning, definition, 4, 5 *see also* motor function
- gagging, 317, 319–320, 321
- gait, 248–277, 360
- analysis of, 145–153*t*, 263, 344, 345–48
 - atypical development, 19, 20, 248–250, 340, 344
 - typical development, 113, 248
- Galant's trunk incurvation, 150*t*
- gastro-oesophageal reflux, 22, 87, 315, 319, 321, 351, 370, 383
- gastrostomy, 173, 315, 320
- General Movements Assessment (GMA), 144–45
- Gillette Functional Assessment Questionnaire (FAQ), 135
- Goal Attainment Scaling (GAS), 70, 141–42
- Goldschmied, E., 29, 46–47, 315
- grasps
- atypical grasps, 295–301
 - developmental levels, 149*t*, 201–2
 - developmental training, 246, 262–63, 288–290, 292, 295–301, 302–3
 - involuntary movement disrupting hand use, 19, 119, 302
- grief, 26–27
- Griffiths Scales, 143–44
- Gross Motor Function Classification System (GMFCS), 11–12, 14, 62, 95, 124
- Gross Motor Function Measure (GMFM), 80–81, 127, 133–34
- Gross Motor Performance Measure (GMPPM), 135
- group work, 79–81, 329–335
- growth, and deformities, 342
- half-kneeling, 169, 189–190, 191, 267, 268–69, 363
- hamstrings, 190–91, 207, 224, 241, 252–53, 345, 347, 349, 357, 363
- hand-arm bimanual intensive therapy, 6
- hands
- assessment of function, 123–24, 133, 137–38, 324–25
 - developmental levels, 184, 186, 201–2, 283*t*–85*t*, 286, 287, 289–290, 300–301, 325
 - developmental training, 115, 166–67, 204, 214–15, 246, 252–53, 261, 262–63, 277–308, 324–26
 - grasping, 19, 119, 149*t*, 201–2, 246, 262–63, 288–290, 292, 295–301, 302–3
 - hand and wrist splints, 295, 297–98, 302, 305–8, 345
 - opening/release, 288, 300–302
 - pointing, 302–3
 - reflexes, 149*t*, 151*t*
 - sign language, 124, 311–12
 - tactile impairments, 3, 154, 298, 299–300
- handwriting, 324–26
- Hanna, S.E., 134
- head
- assessment of, 132–33, 152*t*, 168
 - atypical posture, 242, 365–66, 378
 - developmental training, 174–77, 178–79, 196, 202, 216*f*, 221, 263
 - postural mechanisms, 17, 110, 111, 194, 208, 216*f*, 278, 314, 317–18
- Health and Safety Executive (UK), 388, 389
- health and well-being, 41, 66, 67–69, 85–86, 230
- nutrition, 22–23, 315
 - quality of life (QOL), 62, 68–69, 96, 142–43, 326, 340

- hearing impairments, 20, 330, 333
- hemianopia, 163, 342
- hemiplegia (unilateral cerebral palsy), 15, 124, 240, 259, 298, 339–340, 344–46, 357–58, 359
and sensory impairment, 18, 300
- herbalism, 45
- high stepping, 248, 249
- hip, 17–18
assessment of, 133
bridging, 118, 205
deformities, 223, 224, 241, 338, 345, 346–48, 349–351, 377–78, 381–82
developmental training, 181, 192, 205, 207, 224, 245, 256f
dislocation/subluxation, 192, 338, 349–351, 377, 379, 381–82
pain, 66, 132, 382
and postural control, 162f, 169, 193, 222, 223, 250–51, 255–56, 349–351
- hippotherapy, 85
- hoists, 63, 86, 267, 269, 369, 384–85
- homeopathy, 45
- home visits, 30, 121–22
- horse riding, 70, 85, 230, 360
- hydrotherapy, 85–86
- hyperbaric oxygen therapy, 45
- hyperextension, 176, 191, 241, 265, 296–97, 346, 361–62
- hyperkinesia, 3
- hypersensitivity, 319, 320
- hypertonicity (spastic hypertonus), xiii, 15–18, 72–73, 75, 77, 78, 116–17, 201, 241, 338, 339, 341
assessment of, 129–130
orthoses, 307
and postural control, 213, 232, 240, 245, 250, 261
and selective motor control, 18, 72
spastic diplegia, 8, 62–63, 124, 271, 339–340, 346–48, 357–58
spastic hemiplegia, 344–46, 357–58, 359
spastic tetraplegia (quadriplegia), 124, 339–340, 348, 357–58
use of for compensation, 339
- hypoextensibility, 73
- hypotonicity (hypotonia), 9, 20, 21, 117, 162–63, 200, 338–39
and severe visual impairment, 162
- ice treatment, 74, 351
- immobility, and deformity, 338, 341, 376–77
- impairments, definition, 5
- infants *see* babies (infants, neonates)
- Institute for the Achievement of Human Potential (IAHP), 73, 74
- intellectual impairment, 3–4, 18, 21, 22, 105, 299–300, 332–33
- interdisciplinary teamwork, 42, 55–56, 124–25, 324, 330–32, 370–71
- International Classification of Functioning, Disability and Health (ICF), 4–6, 61, 86, 104, 126, 309, 368
- involuntary movement *see* athetosis; dyskinesia
- joints *see also* casts (plaster casts); orthoses; surgery
assessment of, 131–33, 342–43
joint compression, 214f, 251, 255–56
joint pain, 64
ranges of motion, 131–33
- Kabat, H., 74
- kernicterus, 20
- key points of control, 77
- kicking, 201, 204, 271, 276, 282, 341, 342
developmental levels, 272t–73t, 277t
- KIDSCREEN, 142–43
- kinesio-taping, 86, 297
- knee
assessment of function, 133, 250
deformities, 223, 224, 241, 244, 250, 256f, 339, 346, 347, 361–62
developmental training, 181–82, 189–193, 213, 222, 260f
half-kneeling, 169, 189–190, 191, 267, 268–69, 363
hyperextension, 162f, 260f, 265, 339, 346, 361–62
pain, 66
reflexes, 153t
rotation, 189, 191
splinting, 196, 240, 245, 256f, 263
upright kneeling, 63–64, 169, 171, 190, 192–93, 362
- Knott, M., 74
- kyphosis, 210f, 221, 222, 223–24, 281f, 314, 316, 347, 366, 372, 378
- Lacey Assessment of Preterm Infants, 10
- Landau reflex, 150f
- language acquisition, 310–13 *see also* communication
- lateral sway, 249, 251–52, 255, 261
- lateral weight shift, 249
- 'lead pipe' rigidity, 17, 19–20, 339
- learning cues, 56–58
- learning disabilities, 3–4, 160, 330
- learning styles, 35, 50–51
- learning theory, 2, 25, 80
- legs
deformities, 345, 346–48, 352–57, 376–77
developmental levels, 169–170, 200–201
developmental training, 187–89, 224, 237–249, 250–263
orthoses (splints), 263–68
reflexes, 149t, 150t

- leisure activities, 67–69, 70, 85–86, 230, 277, 326–28, 360
- Lifestyle Assessment Questionnaire - Cerebral Palsy (LAQ-CP), 142
- limbs
- deformities, 344–48, 352–57
 - developmental levels, 169–170
 - developmental training, 187–89
 - orthoses (splints), 263–68, 305–8
 - reflexes, 149*t*, 150*t*
- lordosis, 177, 192, 205, 208, 246, 260*f*, 347, 354*f*, 355*f*, 361, 375
- Lycra suits and splints, 84, 295
- lying *see also* positioning
- night splints, 265, 360
 - prone development, 111, 114–15, 167, 168–193, 204, 334, 350, 393–94
 - prone standers, 238, 245
 - rising from prone, 63–64, 170, 190, 228, 267
 - side lying, 197–200, 202–4, 366
 - supine development, 111, 115, 193–208, 394–95
 - rising from supine, 63–64, 196*f*, 197, 203, 228, 267, 269
 - rolling, 181, 197–200, 206
- magnetic resonance imaging (MRI), 7–8
- malnutrition, 22–23, 315
- Manual Ability Classification System (MACS), 12–13, 14
- manual handling *see* moving and handling
- Martin, J.P., 105, 106, 107, 249
- massage, 45, 46
- McLellan, D.L., 51, 95, 116
- Measure of Processes of Care (MPOC), 29, 38
- medication, 17, 86–87, 116
- Melbourne Assessment of Unilateral Upper Limb Function (MA), 137–38
- memory, 3, 18, 326
- 'mermaid crawl,' 55, 180, 341
- meta-analyses, 97
- methodology (research methodology), 94–95, 96–102
- mindfulness, 41, 69
- mirror movements, 283*t*, 298
- mirrors, 57, 215*f*, 244, 304
- mistakes, 50–51, 52, 60, 75
- mobility aids, 6, 383–88
- powered mobility aids, 383–84, 386–88
 - tricycles, 192, 205, 228, 263, 277
 - walking aids, 115, 168, 256*f*, 257*f*, 258–263, 383, 384–85
 - wheelchairs, 62, 66, 68–69, 136, 148, 369, 378, 383–84, 385–88
- Mobility Opportunities Via Education (MOVE programme), 88–89
- Modified Ashworth Scale (MAS), 129, 130
- Modified Tardieu Scale, 129–130
- Moro reaction, 21, 150*t*, 201–2, 227
- mothers, 37*f*, 165*f*, 240*f see also* parents
- motivation and rewards, 2, 26, 30, 33–34, 39–40, 50–54, 58–59, 164, 203–4, 311, 313
- for adolescents, 65
 - in group work, 334–35
- motor function, 115
- activation patterns, 17–18, 77, 78–79, 110, 159, 232, 340–41, 352, 356*f*
 - in adolescents and adults, 62–63, 67–69
 - assessment of *see* assessment
 - atypical synergies, 119–120, 127, 341
 - child's own strategies, 50, 54–55, 206, 207*f*
 - co-contraction, 18, 78, 106, 116, 339, 340–41
 - in daily life *see* daily life
 - developmental levels *see* developmental levels
 - and fatigue, 19, 22, 52, 63, 66–67
 - goal-directed movements, 53, 159
 - learning methods, 49–60, 75, 88–89, 108–10, 157–58, 300, 344, 352, 360
 - cues for learning, 56–58, 164–68
 - practice, 59–60, 117–120
 - measurement tools, 125–145
 - movement patterns, 74, 110, 117–18, 187–89, 197–200, 205, 227–28, 252–54, 280–82, 290–300, 342
 - postural control *see* postural control
 - progressive pattern movements, 73–74
 - selective motor control, 18, 72, 117–18, 126–27, 250
 - visual-spatial competence, 3
 - voluntary motion, 108–10, 119–120
- motor growth curves, 134
- motor training *see* developmental training
- moving and handling (manual handling), 63, 369, 381, 388–390
- MRI (magnetic resonance imaging), 7–8
- multiculturalism, 29–30 *see also* culturally appropriate therapy
- muscle education, 71–73
- musculoskeletal deformities *see* deformities
- Muslim families, 30
- myositis ossificans, 353
- nasogastric tubes, 320
- National Institute for Health and Care Excellence (NICE), xiv, 10, 45, 81, 97, 236, 307, 379–380, 385
- neck, 150*t*, 151*t*, 341, 365–66
- negative supporting reaction, 151*t*
- neonates *see* babies (infants, neonates)
- neurodevelopmental treatment (NDT), 50, 75–77

- neurofacilitation treatments, 94, 104, 118
- neuroimaging, 7–8
- neuromotor development, 75
- neuromuscular electrical stimulation, 83–84
- neuronal group selection theory, 90
- neuroplasticity, 89–90
- night splints, 265, 360
- non-verbal communication, 3, 32, 52, 124, 311, 313
- Novak, I., xi
- nutrition, 22–23, 315
- nystagmus, 21
- occupational therapy, 29, 81, 309, 314–15, 324–25, 326
 and alternative treatments, 44
 assessment, 122, 140–41, 156, 368–69
 and botulinum toxin A, 358–59
 group work, 330–31
- older people *see* adults
- one-minute walk test, 136
- ontogenetic development, 73, 78, 111
- optical reflex, 152*t*
- oromotor dysfunction, 322
- orthoses, 182, 243*f*, 245, 250, 263–68
 abduction splints, 203, 224, 245, 263, 350
 ankle-foot orthoses, 68*f*, 250, 251, 259, 264–66, 345, 347, 351, 362, 363
 body orthoses, 366, 377
 braces and calipers, 72
 elbow orthosis, 181, 256*f*, 257*f*, 261
 hand and wrist splints, 295, 297–98, 302, 305–8, 345
 kinesio-taping, 86, 297
 knee splints, 196, 240, 245, 256*f*, 263, 361
 Lycra suits and splints, 84, 295
 night splints, 265, 360
- osteoarthritis, 64, 66
- osteopathy, 45
- Paediatric Reach Test, 135
- pain, 22, 64, 66, 87, 342, 375, 382, 383
- palmar flexion, 295, 306*f*
- parachute reactions, 107, 170, 194, 241
- parents, 25–48
 evaluation of services, 28–29, 37–38, 80
 family-centred services, xii, 25–30, 46–48
 health and well-being, 22, 26–28, 35–36, 41–42, 315, 318, 351
 learning styles, 35
 parent-child relationship, 26–28, 36–37, 38–41, 46–48, 60, 163–64, 318–320
 participation in goal-setting, 26, 30–32
 positioning, 159, 204, 237–38, 314, 317–18, 349–352, 361, 363, 370
 relationship with adolescents, 65
 relationship with therapist, 31–32, 41, 42, 75, 348, 353
 support for, 26–28, 41–42, 315, 318, 348, 351
 and task analysis, 32–34
- participant observation, definition, 102
- participation, definition, 4, 5
- passive observation, definition, 102
- Patient Perspective on Care and Rehabilitation (POCR), 27, 38
- patterning therapy (progressive pattern movements), 73–74
- Peabody Developmental Motor Scales, 143
- Pediatric Evaluation of Disability Inventory (PEDI), 138–39
- Pediatric Quality of Life Inventory (PedsQL), 142
- pelvis
 deformities, 223, 224, 241, 338, 345, 346–48, 377–78
 developmental levels, 162*f*, 169, 193–94, 202, 208–9, 222
 developmental training, 181, 197–98, 205, 223, 231, 245
 and postural control, 162*f*, 169, 192, 222, 223–24, 250–51, 255–56, 349–351, 374–78
- perceptual development, 304–5
- perceptual impairments, 3, 18, 105, 110, 158, 161–68, 277, 299–300, 302
- periventricular leukomalacia, 8
- personality, 5–6, 20, 333
- Pető approach, 79, 119, 259
- Phelps, W.M., 71–73
- physiotherapy, 29, 36, 95–96, 309, 314–15, 326
 for adolescents, 66, 160
 and alternative treatments, 44
 assessment, 155–56
 and botulinum toxin A, 359–360
 group work, 330–31
 postoperative therapy, 87, 112, 132, 340, 347, 348–49, 351, 357, 361
 and sensory issues, 82
- Physiotherapy Evidence Database (PEDro), 97, 98, 100
- pivoting, 273
- pivot prone, 78, 169, 186, 187
- plantarflexion, 215, 264–65, 340, 362, 363
- plasters (casts), 99, 245, 264–65, 351, 352–57, 359, 362–63
- play, 6, 54, 163, 165, 166–67, 183, 230, 289, 326–28, 334–35
 developmental levels, 327–28
 playgroups, 329–330
 practical suggestions, 328
 treatment during, 36–37
- pointing, 302–3

- positioning, 36–37, 39, 56–58, 159, 349–351, 362, 366, 371–386
 during feeding, 317–18, 370, 375
 postoperative therapy, 87, 112, 132, 340, 347, 348–49, 351, 357, 361
 postural adjustment *see* counterpoising (postural adjustment)
 postural control, 18, 19, 21, 75 *see also* equipment arms, 278–282
 assessment of, 123, 271
 atypical posture *see* atypical postures
 developmental training, 110–16, 159, 186–89, 224, 242–48, 250–263, 267–69, 270f, 271–77, 363, 364f
 head and trunk, 17, 168–69, 193–94, 196, 208, 248–49, 278–79, 314, 317–18, 377, 378
 horse riding, 85, 230, 360
 and hypertonicity, 16–17, 19–20, 250, 261
 in play, 36–37
 in postoperative care, 349–351
 and voluntary motion, 109–10
 postural mechanisms, 105–8, 167–68, 208, 212–16, 223–25, 236, 339, 343, 344
 antigravity mechanism, 21, 106, 128, 230, 233f, 237, 245, 341, 346
 counterpoising, 106, 137, 168f, 169–170, 187–89, 193–94, 208, 209, 227f, 231–32, 241, 252–55, 276–77, 279f
 equilibrium reactions, 107–8, 111
 and hypotonia, 162–63
 righting (rising) reactions, 79, 106–7, 115, 179–180, 193, 194, 209, 228–230, 232, 267–69, 279–280
 saving reactions, 107, 128, 153t, 170–71, 184, 191, 194, 210, 211f, 232–33, 240–41, 249–250, 271, 280
 tilt reaction, 36f, 107–8, 120, 152t, 170, 171, 191, 194, 209–10f, 211f, 212f, 230, 232, 240–41, 249, 273–77, 396
 powered mobility aids, 383–84, 386–88
 Prechtel, H.F.R., 144–45
 Preferences for Activities of Children (PAC), 140
 preterm babies, 10–11
 productivity (school/work), 309, 324
 progressive pattern movements (patterning therapy), 73–74
 progressive strength training (PST), 82–83, 111, 340, 351
 prone development, 111, 114–15, 167, 168–193, 204, 334, 350, 393–94
 prone standers, 238, 245
 rising from prone, 63–64, 170, 190, 228, 267
 proprioception, 105, 165–66, 294, 353
 proprioceptive neuromuscular facilitations (PNF), 74–75
 quadriplegia, 14, 16f, 18, 124, 222, 316–17, 339–340, 348 *see also* bilateral cerebral palsy
 qualitative research, validity and reliability, 95, 96–97, 98–99
 quality of life (QOL), 62, 67–69, 142–43, 326, 340
 health-related (HRQOL), 67–68, 96, 142
 Quality of Upper Extremity Skills Test (QUEST), 137
 quantitative research, validity and reliability, 96, 98–102
 randomised controlled trial, definition, 98
 reaching, 290–94
 rebound therapy, 86
 reciprocation, 72
 record-keeping, 155–56
 reflective learning, 51
 reflex creeping, 78–79
 reflexes, 74, 78, 149t–153t, 270–271
 in babies, 21–22, 197–98, 201–2, 204, 235, 237, 287, 315, 344
 tonic reflexes, 21, 76, 150t–51t, 201–2, 341
 reflexology, 45
 reflux, 22, 87, 315, 319, 321, 351, 370, 383
 relaxation techniques, 41, 44–45, 72, 74
 research studies, validity and reliability, 10, 29, 93–102, 141
 resistance, 74, 78–79, 352
 for improving stability, 79, 184, 189, 247, 251, 256
 interactions between body parts, 119, 180
 for strengthening, 75, 82
 respiratory problems, 18, 65, 68, 114, 173, 366
 rewards *see* motivation and rewards
 righting (rising) reactions, 79, 106–7, 179–180, 193, 194, 209, 228–230, 232, 267–69, 279–280
 rising, 63–64, 79
 developmental levels, 106–7, 115, 170, 172, 232
 developmental training, 179–180, 181, 193, 194, 196–97, 200, 228–230, 267–69, 279–280
 rocking *see* anteroposterior weight shift (rocking)
 Rogers, C.R., 25
 rolling, 172, 174, 181, 186–87, 197–200, 206, 207, 235
 Rood, M.S., 73, 77–78, 111
 rooting, 149t, 315
 rotation, 119, 189, 191, 197–98, 257f, 352
 Royal College of Occupational Therapists (UK), 81, 309
 Royal National Institute of Blind People (UK), 162, 168
 running, 248, 249, 255, 262, 273
 sacral sitting, 221, 222, 375
 sampling techniques, 98–99, 100–101, 102

- saving reactions, 107, 128, 153*t*, 170–71, 184, 191, 194, 210, 211*f*, 233–35, 240–41, 249–250, 271, 280
- school, 30, 318, 324–26, 383
- scissoring, 237, 245, 248, 357, 384
- scoliosis, 65, 120, 194, 203*f*, 222, 223, 224, 226*f*, 240, 341, 342, 343, 347, 350, 355*f*, 365–66, 371, 378, 381–82
- Scottish Centre for Children with Motor Impairments, 81
- Scottish Intercollegiate Guidelines Network (SIGN), 97
- seating *see* chairs/seating systems
- seesaw game, 36*f*
- Selective Control Assessment of Lower Extremity (SCALE), 127
- selective dorsal rhizotomy, 17, 87, 116
- selective motor control, 18, 72, 117–18
- self-care tasks, 62, 63, 69, 160, 164*f*, 313–324, 335, 368–370, 378
- self-report measures, 140, 142–43
- sensory integration, 81–82
- sensory stimulation, 10, 74, 77–78, 118–19, 163
- sensory systems, 105–6
 - assessment of, 124, 154
 - hand-eye coordination, 283*t*–85*t*, 286–298, 300, 303, 304–8
 - hypersensitivity, 319, 320
 - sensory impairments, 3, 18, 20, 298, 299–300, 302, 319, 330, 333, 338, 342, 353
- shoes (boots, trainers), 265–66, 363
- shoulder, 18, 119
 - assessment of, 128, 133
 - developmental levels, 168*f*, 169, 208
 - developmental training, 181, 197–99, 314
- showering, 368–69, 378
- Shriners Hospital for Children Upper Extremity Evaluation (SHUEE), 138
- siblings, 43, 65
- side lying, 197–200, 202–4, 366
- side sitting, 224, 230
- sign language, 124, 311–12
- single-event multilevel surgery (SEMLS), 345, 347
- single parents, 28
- sinking posture, 255, 256*f*
- sitting, 111, 212–16, 334, 395–96
 - assessment of, 135, 150*t*
 - atypical posture, 220–23, 225, 227, 375
 - developmental training, 191, 196*f*, 197, 203, 208–30, 267–69, 314–15
 - orthoses, 366
 - seating systems, 205, 215, 217–19, 221, 223–25, 315, 369, 370, 371–79
 - side sitting, 224, 230
- Sitting Assessment for Children with Neuromotor Dysfunction (SACND), 135
- six-minute walk test, 136
- sleep, 265, 351–52, 360, 366, 381–83
 - problems in child, 20, 41
 - problems in parents, 22
- social development, 3–4, 22, 70, 318–320, 333, 334, 368–69
- spasticity (spastic hypertonus), xiii, 15–18, 72–73, 75, 77, 78, 116–17, 201, 241, 338, 341
 - assessment of, 129–130
 - orthoses, 307
 - and postural control, 213, 232, 239, 245, 250, 261
 - and selective motor control, 18, 72, 339
 - spastic diplegia, 8, 62–63, 124, 271, 339–340, 346–48, 357–58
 - spastic hemiplegia, 344–46, 357–58, 359
 - spastic tetraplegia (quadriplegia), 124, 339–340, 348, 357–58
 - use for compensation, 339
- speech and language therapy, 158, 311–13, 319, 322, 330–31
- splinting, 84, 182, 243*f*, 245, 363 *see also* casts (plaster casts); orthoses
 - abduction splints, 203, 224, 245, 263, 350
 - elbow splints (gaiters), 256*f*, 257*f*, 261
 - hand and wrist, 295, 297–98, 302, 305–8, 345
 - knee, 196, 240, 245, 256*f*, 263, 361–62
 - night splints, 265, 360
- spondylosis, 64
- spoon feeding, 320–21
- sport and leisure activities, 67–69, 70, 85–86, 230, 277, 326–28, 360
- stair banisters, 392
- stair climbing, 136, 168, 191, 192, 253*f*, 265, 271–73, 363
- standing, 111, 115, 151*t*, 153*t*, 334, 349, 396
 - atypical posture, 238–242, 339
 - developmental training, 230–248, 267–69, 271, 314, 363, 364*f*
 - standing frames, 50*f*, 73, 114, 236, 244, 245–46, 379–381
- startle reflex, 150*t*, 287
- status dystonicus, 20
- stepping, 115
- stereotyping, 30
- stiffness, 15
- strength training, 63, 73, 82–86, 108–9, 111, 116–17, 269, 277, 340, 344, 360
- stretching, 191, 252, 363, 365, 366
 - corrective casts, 353, 359, 360
 - leisure activities, 86, 360
 - postoperative therapy, 347, 351
 - prolonged stretch, 73, 351–52, 353, 360, 364

- subluxation, 192, 338–39, 343–44, 349–351, 377, 379, 381–82
 sucking, 149*t*
 supine development, 111, 115, 193–208, 394–95
 rising from supine, 63–64, 196*f*, 197, 203, 228, 267, 269
 rolling, 181, 197–200, 206
 support groups, 27
 supporting reaction (antigravity mechanism), 21, 106, 128, 230, 233*f*, 237, 245, 341, 346
 support services, 64–65, 399
 suprasternal palsy, 18
 surgery, 17, 87, 95–96, 116, 266, 315, 347, 348–351, 353, 366
 orthopaedic, 345–46, 349, 351
 Surveillance of Cerebral Palsy in Europe, 7
 swallowing problems (dysphagia), 67, 315, 317
 sway, 248, 249, 251–52, 255, 261
 swimming, 30, 70, 85–86
 symmetrical tonic neck reflex, 151*t*
 synergies, atypical development of, 72, 119–120, 127, 341
 systematic reviews, validity and reliability, 97, 141
 systems-based task-oriented approach, 87–88

 tables, 321, 371
 tactile impairments, 3, 154, 298, 299–300, 353
 tailor sitting, 221, 225, 323*f*, 361
 taping, 86, 297
 Tardieu Scale, 129–130
 targeted training, 83
 task analysis, 32–34, 55–56, 80, 88–89, 121–22, 392
 teamwork, 4, 27, 42–43, 55–56, 124–25, 324, 330–32, 370–71
 multidisciplinary team (MDT), 42, 313, 324
 Test of Infant Motor Performance (TIMP), 145
 tetraplegia, 14, 124, 339–340, 348, 357–58 *see also*
 bilateral cerebral palsy
 theories, 81, 89–92, 94, 104
 therapists
 confidentiality, 61, 68
 professional development, 29, 51
 relationship with adolescents, 65–66
 relationship with parents, 31–32, 41, 42, 75, 348, 353
 support for, 27, 399
 teamwork, 4, 27, 42–43, 55–56, 64–65, 124–25, 324, 330–32, 370–71
 therapy goals, xiii–xiv, 2, 26, 30–32, 53, 62, 65, 122, 141–42, 156, 343–44, 357–58, 359–360
 thumbs, 297–98, 306*f*
 tilt reaction, 36*f*, 107–8, 120, 152*t*, 170, 171, 191, 194, 209–10*f*, 211*f*, 212*f*, 230, 232, 240–41, 249, 273–77, 353, 396
 timed sit-to-stand test, 136
 timed stairs test, 136
 timed up and go (TUG), 136
 toe clenching, 241
 toe walking, 255, 345, 346, 347, 356
 toilet seats, 369
 tone inhibitory casts, 352–57
 tonic reflexes, 21, 76, 150*t*–51*t*, 201–2, 341
 topographical classifications, 3, 14–15
 torticollis, 204, 222, 341
 'traffic light' evaluation, x, 77
 trainers, 266, 363
 trampolining, 86
 transdisciplinary teamwork, 42
 traumatic brain injury, 7, 160
 treadmill training, 84–85
 treatment goals *see* therapy goals
 tremor, 20
 tricycles, 192, 205, 228, 263, 277
 trunk
 deformities, 365–66
 developmental levels, 168, 196
 orthoses, 366, 377
 postural mechanisms, 17, 108, 110, 193–94, 208, 221, 224, 245, 263, 278–79, 314, 317–18
 reflexes, 150*t*, 152*t*
 segmental assessment of trunk control (SATCo), 83

 ulnar deviation, 302, 344–45
 unilateral cerebral palsy, 15, 124, 240–241, 259, 298, 339–340, 344–46, 357–58, 359
 and sensory impairment, 18, 300
 universal design, 391–92
 upright kneeling, 63–64, 169, 171, 190, 192–93, 362
 urinary problems, 67

 valgus feet, 223, 238, 241, 266, 345, 346–47, 351, 363–64, 365*f*
 valgus knees, 244, 339, 347
 varus feet, 266, 346, 351, 352, 364
 varus knees (bow legs), 339
 verbal guidance, 56, 58
 vestibular perception, 105, 165–66
 visual feedback, 57–58
 visual impairments, 3, 18, 23, 105, 158, 161–68, 299–300, 342
 and hand function, 166–67, 229–230, 294, 302
 hypotonia and severe visual impairment, 162
 and language development, 166
 and postural control, 108, 110, 115, 162
 residual vision, 166
 visual perception, 105–6
 Vojta, V., 78–79, 111, 119, 180
 voluntary motion, 108–10, 119–120
 Voss, D.E., 74

- waddling, 248, 255, 348
- walking, 17–18, 111, 124, 167–68, 396
 - assessment of, 145–153*t*, 233–36, 344, 345–48
 - bear-walk, 170, 172*f*, 189, 190–91
 - changes in function, 9–10, 62–63, 66–67, 69
 - developmental training, 190–91, 247–263, 271–77
 - prognosis for, 62–63, 235–36
 - speed, 148, 247, 255, 262
- walking aids, 115, 168, 236–37, 256*f*, 257*f*, 258–263, 383, 384–85
- weakness, 19, 64, 87, 113–14, 116–17, 127–29, 339–340
- wedges, 176–78, 205, 224, 225, 350
- WeeFIM (Functional Independence Measure for Children), 139
- weight bearing
 - postoperative therapy, 349
 - in prone development, 176–78, 181–82, 184, 186, 191
 - in sitting, 216*f*, 221–22
 - standing and walking, 237–38, 242–44, 245, 251–52, 253, 257*f*, 263, 353, 364*f*
- weight shift
 - anteroposterior (rocking), 169, 232, 247*f*, 249, 252*f*
 - lateral, 249
- weight training, 82–83
- wheelchairs, 62, 66, 68–69, 369, 378, 385
 - powered wheelchairs, 383–84, 386–88
 - wheelchair skills, 136, 148, 386–88, 397
- windswept posture, 201, 203*f*, 229–230, 341, 350–51, 365–66, 377
- World Health Organization (WHO), 4, 309, 368
- Worster-Drought Syndrome, 18
- wrist splints, 295, 297–98, 302, 345
- W-sitting, 220*f*, 221, 222
- Wu, Y.W., 235